Meet ISMP Guidelines

for Safe Medication Use in Perioperative and Procedural Settings by 2025 with SAFE Label System



Harmful medication errors are a significant public health problem, causing at least 1 death every day and injuring 1.3 million people annually.¹

In the perioperative setting, medication errors occur frequently in all phases of perioperative care and are a common cause of morbidity and mortality.²⁻⁴

Overall, medication errors are estimated to occur in at least **1 in every 133** doses administered during anesthesia alone.⁵⁻⁶

Codonics Safe Label System addresses these risks and introduces consistent Best Practices that create standardization and bring barcode medication administration (BCMA) and long overdue electronic safety checks that improve patient care while improving the anesthesia workflow. It's helping to meet these new ISMP guidelines⁷, several of which are targeted for implementation by 2025.





To help improve medication safety,

ISMP worked with clinical experts, professional organizations and industry leaders on best practice guidelines across all phases of perioperative care. These **2022 Guidelines** have been made available to hospitals, ambulatory surgery centers and other procedural locations to address identified national gaps in perioperative and procedural medication safety and provide a stepping stone and support for further implementation to reduce harmful patient events.



The need to improve medication safety in the OR has received increasing attention in recent years from American Society of Health-System Pharmacists (ASHP), the Anesthesia Patient Safety Foundation (APSF), the Association of periOperative Registered Nurses (AORN) and the Institute for Safe Medication Practices (ISMP). The Joint Commission (TJC) continues to include perioperative medication safety practices as part of its National Patient Safety Goals (NPSG).

"Bar code medication administration has been the standard of care in the majority of hospital locations for nearly two decades. Finally, the Institute for Safe Medication Practices has recently advocated for bringing that same standard of care to our anesthesia practice."

--- Joyce A. Wahr, MD, FAHA

Vice Chair, Quality and Safety, Department of Anesthesiology, University of Minnesota

AT PREPARATION		
ISMP Guideline 4.3	Eliminate the use of handwritten labels in perioperative/procedural areas by 2025 .	✓
ISMP Guideline 4.4	Include a machine-readable code (e.g. barcode, radio frequency identification [RFID]) on all syringe and infusion labels, including those that are PRACTITIONER-PREPARED , by 2025.	~
ISMP Guideline 4.5	Label PRACTITIONER-PREPARED syringes of medications with, at the minimum, the full name, concentration/dose of the drug, name or initials of the preparing practitioner, as well as an expiration date (when not used in 24 hours) and time (if expiration occurs in less than 24 hours). Application of an anesthesia color-coded drug class label alone is not sufficient.	✓




	AT ADMINISTRATION	
ISMP Guideline 10.11	Use machine-readable coding (e.g. BARCODE SCANNING , RFID) in pre operative/ pre procedural <u>and</u> post operative/ post procedural settings to verify patients and medications/solutions prior to administration.	√
ISMP Guideline 10.12	Take steps to implement machine-readable coding (e.g. BARCODE SCANNING , RFID) in intra operative/ intra procedural workflows to confirm medication/solution selection prior to administration.	V
ISMP Guideline 10.13	Take steps to implement and integrate machine-readable coding (e.g. BARCODE SCANNING , RFID) to support real-time EHR documentation of medication doses and fluid administration in all preoperative/preprocedural, intraoperative/intraprocedural and postoperative/postprocedural settings.8 *Future when integrated with BD's Intelliport System	~



Additionally, AT ADMINISTRATION		
Meets ISMP Guideline 6.6	Take steps toward the implementation of bidirectional (i.e. autoprogramming and auto-documentation) SMART INFUSION PUMP interoperability with the EHR in all preoperative/preprocedural, intraoperative/intraprocedural, and postoperative/postprocedural settings. ⁹	~
ISMP Targeted Medication Safety Best Practices for Hospitals 2022/2023 Best Practice 18	Maximize the use of barcode verification prior to medication and vaccine administration by expanding use beyond inpatient care areas. ¹⁰	✓







on preventing diversion of controlled substances¹¹





Call Codonics today at 800.444.1198 or visit www.codonics.com for more information.

References:

- | World Health Organization, WHO lounches global effort to halve medication-related errors in 5 years, Available at: wave immp. org.ext/279. Published 2017. Accessed July 30, 2018.

 2 Beyes SC, Hoka RW, Bedark SC, Medication errors in 16 o. St.—a secondary analysis of Medinors. AIDRN 1, 2003;7(11);123;1259-1322.4.

 2 Beyes SC, Hoka RW, Bedark SC, Medication errors in 16 o. St.—a secondary analysis of Medinors. AIDRN 1, 2003;7(11);123;1259-1322.4.

 3 Boylin JL (Link Richard Cardior Cardior School Cardior C



17991 Englewood Drive Middleburg Heights, OH 44130 USA

- +1.440.243.1198
- +1.440.243.1334 Fax Email info@codonics.com www.codonics.com